

## Dual enzymatic activity of the pathogenesis-related protein TcPR-4 from *Theobroma cacao*: ribonuclease and $\text{Ca}^{+2}$ and $\text{Mg}^{+2}$ dependent deoxyribonuclease activities

Menezes, SP<sup>1</sup>; Silva, EMA<sup>1</sup>; Lima, EM<sup>1</sup>; Gramacho, KP<sup>2</sup>; Pirovani, CP<sup>1</sup>; Gesteira, AS<sup>3</sup>; Micheli, F<sup>1,4</sup>.

<sup>1</sup>Universidade Estadual de Santa Cruz, Centro de Biotecnologia e Genética, Rodovia Ilhéus-Itabuna, Km 16, 45662-000, Ilhéus-BA, Brasil; <sup>2</sup>CEPLAC/CEPEC, 45600-970, Itabuna-BA, Brasil; <sup>3</sup>Embrapa Mandioca e Fruticultura, Rua Embrapa, s/nº, 44380-000, Cruz das Almas-BA, Brasil; <sup>4</sup>CIRAD, UMAR AGAP, Avenue Agropolis, Montpellier, France.

**Keywords:** Barwin domain, nuclease, *Moniliophthora perniciosa*.

The class 4 pathogenesis-related proteins (PR4) are classified as chitinases and contain a conserved Barwin domain. The *TcPR-4b* cDNA identified from a library of *Theobroma cacao* L. pod (genotype TSH1188) infected by *Moniliophthora perniciosa* also presents the Barwin domain with six conserved cysteine residues, but lacks the chitin-binding site and for this reason was classified as class II PR4. The *TcPR-4b* gene was cloned into pET28a and the resulting in frame fusion plasmid was used to transform *Escherichia coli* Roseta (DE3) for protein expression. The expression of the TcPR-4b recombinant protein was induced by 0.4 mM isopropyl-β-D-thio-galactoside and purified by immobilized metal affinity chromatography with TALON<sup>®</sup> Metal Affinity Resin. To determine the DNase activity of the purified recombinant TcPR-4b protein, 1 µg of purified pGEM-T<sup>®</sup> Easy Vector DNA was incubated with different protein amounts (2, 4, 6, 8, 10, 12, 14, 16, 18 and 20 µg) in the presence or absence of 10 mM of  $\text{MgCl}_2$  or 1 mM of  $\text{CaCl}_2$  overnight at room temperature. RNase activity of recombinant TcPR-4b was performed using different protein amounts (5, 10, 15, 20 and 25 µg) incubated for 30 min with 5 µg of RNA extracted from *Solanum lycopersicum* leaves. The reaction products were analyzed in 1.5% agarose electrophoresis gel. The TcPR-4b protein recombinant showed both DNase and RNase activity. DNase activity was observed only in the presence of  $\text{Mg}^{+2}$  and  $\text{Ca}^{+2}$  ions. The results of this study suggest that TcPR-4b may act as nuclease during the infection of cacao plants with *M. perniciosa*. Financial Support: CNPq, BNB, FINEP/RENORBIO, CAPES.